# BULLITT COUNTY REPORT OF ENDANGERED, THREATENED, AND SPECIAL CONCERN PLANTS, ANIMALS, AND NATURAL COMMUNITIES OF KENTUCKY

PRESERVES COMMISSION 801 SCHENKEL LANE FRANKFORT, KY 40601 (502) 573-2886 (phone) (502) 573-2355 (fax)

www.naturepreserves.ky.gov

# Kentucky State Nature Preserves Commission Key for County List Report

Within a county, elements are arranged first by taxonomic complexity (plants first, natural communities last), and second by scientific name. A key to status, ranks, and count data fields follows.

### **STATUS**

KSNPC: Kentucky State Nature Preserves Commission status:

USESA: U.S. Fish and Wildlife Service status:

SOMC = Species of Management Concern

## **RANKS**

GRANK: Estimate of element abundance on a global scale:

G1 = Critically imperiled GU = Unrankable

G2 = Imperiled G#? = Inexact rank (e.g. G2?)
G3 = Vulnerable G#Q = Questionable taxonomy

G4 = Apparently secure G#T# = Infraspecific taxa (Subspecies and variety abundances are coded with a 'T' suffix; the 'G'

G5 = Secure portion of the rank then refers to the entire species)

GH = Historic, possibly extinct GNR = Unranked GX = Presumed extinct GNA = Not applicable

SRANK: Estimate of element abundance in Kentucky:

S1 = Critically imperiled SU = Unrankable Migratory species may have separate ranks for different

S2 = Imperiled S#? = Inexact rank (e.g. G2?) population segments (e.g. S1B, S2N, S4M):

S3 = Vulnerable S#Q = Questionable taxonomy S#B = Rank of breeding population
S4 = Apparently secure S#T# = Infraspecific taxa S#N = Rank of non-breeding population
S5 = Secure SNR = Unranked S#M = Rank of transient population

SH = Historic, possibly extirpated SNA = Not applicable

SX = Presumed extirpated

### **COUNT DATA FIELDS**

# OF OCCURRENCES: Number of occurrences of a particular element from a county. Column headings are as follows:

- E currently reported from the county
- H reported from the county but not seen for at least 20 years
- F reported from county & cannot be relocated but for which further inventory is needed
- X known to be extirpated from the county
- U reported from a county but cannot be mapped to a quadrangle or exact location.

The data from which the county report is generated is continually updated. The date on which the report was created is in the report footer. Contact KSNPC for a current copy of the report.

Please note that the quantity and quality of data collected by the Kentucky Natural Heritage Program are dependent on the research and observations of many individuals and organizations. In most cases, this information is not the result of comprehensive or site-specific field surveys; many natural areas in Kentucky have never been thoroughly surveyed, and new species of plants and animals are still being discovered. For these reasons, the Kentucky Natural Heritage Program cannot provide a definitive statement on the presence, absence, or condition of biological elements in any part of Kentucky. Heritage reports summarize the existing information known to the Kentucky Natural Heritage Program at the time of the request regarding the biological elements or locations in question. They should never be regarded as final statements on the elements or areas being considered, nor should they be substituted for on-site surveys required for environmental assessments.

KSNPC appreciates the submission of any endangered species data for Kentucky from field observations. For information on data reporting or other data services provided by KSNPC, please contact the Data Manager at:

Kentucky State Nature Preserves Commission 801 Schenkel Lane Frankfort, KY 40601 phone: (502) 573-2886 fax: (502) 573-2355

email: naturepreserves@ky.gov internet: www.naturepreserves.ky.gov

County	/ Taxonomic Group	Scientific name	Common name	Statuses	Ranks		# of	Occ	urren	ices
	Habitat					E	Н	F	X	U
Bullitt	Vascular Plants Rich, moist woods, thickets and we	Agrimonia gryposepala oodland borders.	Tall Hairy Groovebur	Т/	G5 / S1S2	0	1	0	0	0
Bullitt	Vascular Plants CEDAR GLADES AND PRAIRIES	Carex crawei , ALSO REPORTED IN CALCAREOUS SHORES AND ME	Crawe's Sedge ADOWS.	S/	G5 / S2S3	2	0	0	0	0
Bullitt	Vascular Plants Calcareous bluffs and rocks (Gleat	Cheilanthes feei son & Cronquist 1991)	Fee's Lipfern	E/	G5 / S1	1	0	0	0	0
Bullitt	Vascular Plants Reported in savannahs, bogs, mea	Fimbristylis puberula adows and prairies, open limestone, chert or sandstone glad	Hairy Fimbristylis des; cedar glades on limestone in KY.	Т/	G5 / S2	1	0	0	0	0
Bullitt	Vascular Plants Wet places in cedar glades.	Juncus filipendulus	Ringseed Rush	Т/	G5 / S2?	1	0	1	0	0
Bullitt	Vascular Plants In full sun on flat-bedded outcrops	Leavenworthia exigua var. laciniata of silurian limestone or dolomite in shallow soils of glades,	Kentucky Gladecress rock oucrops, pastures and lawns.	E/ SOMC	G4T1T2 / S1S2	44	0	8	15	0
Bullitt	Vascular Plants Dry woods, barrens, and prairies,	Oenothera triloba often calcareous; in KY, glades, dry limestone soil, rock out	Stemless Evening-primrose crops in fields.	Т/	G4 / S1S2	1	0	0	1	0
Bullitt	Vascular Plants Low grounds, prairies, and rich wo	Perideridia americana ods.	Eastern Yampah	Т/	G4 / S2	1	0	0	0	0
Bullitt	Vascular Plants Calcareous soil in prairies, and gla	Spiranthes magnicamporum ndes.	Great Plains Ladies'-tresses	Т/	G4 / S2	3	0	0	0	0
Bullitt	Vascular Plants Damp (although sometimes seaso	Spiranthes ochroleuca nally only) acid soil of open woods and grassy openings.	Yellow Nodding Ladies'-tresses	Т/	G4 / S2?	0	1	0	0	0
Bullitt	Vascular Plants Dry open ground, prairies, glades	Sporobolus heterolepis and woodland openings near glades, rocky cliffs.	Northern Dropseed	E/	G5 / S1	1	0	0	0	0
Bullitt	Vascular Plants Open dry woods, bluffs and prairie	Symphyotrichum pratense s. Occurs with prairie vegetation and in cedar glades in KY.	Barrens Silky Aster	S/	GNR / S3	1	0	0	0	0
Bullitt	Vascular Plants CALCAREOUS BARRENS, GLAD	Viola septemloba var. egglestonii DES AND DRY PRAIRIES ON SILURIAN AND MISSISSIPP	Eggleston's Violet IAN LIMESTONES.	S/	G4 / S3	20	3	0	2	0
Bullitt	Freshwater Mussels LARGE RIVER SPECIES THAT IN STANSBERY 1976).	<i>Obovaria retusa</i> NHABITS GRAVEL AND SAND BARS (BOGAN AND PARM	Ring Pink MALEE 1983, GOODRICH AND VAN DER SCI	E / LE HALIE 1944, NEE	G1 / S1 L AND ALLEN 1964,	0	0	0	1	0
Bullitt	Freshwater Mussels USUALLY FOUND IN LARGE RIV	Plethobasus cooperianus /ERS IN SAND AND GRAVEL SUBSTRATES (AHLSTEDT	Orangefoot Pimpleback 1983, BOGAN AND PARMALEE 1983, MILLE	E / LE :R, A.C. ET AL. 19	G1 / S1 986).	0	0	0	1	0
Bullitt	Freshwater Mussels Usually found in large rivers in cur	Plethobasus cyphyus rent on mud, sand, or gravel bottoms at depth of 1-2 meters	Sheepnose or more (Baker 1928, Parmalee 1967, Gordon	E / C n and Layzer 1989	G3 / S1 9).	1	0	0	0	0
Bullitt	•	Pleurobema clava nall streams and rivers (Goodrich and Van Der Schalie 1944 te and consequently difficult to find (Watters 1987).	Clubshell ; Ortmann 1919,1925), although in Kentucky it	E / LE is known from mo	G2 / S1 oderately large rivers.	0	0	0	1	0
Bullitt		Simpsonaias ambigua STRATE SUCH AS SOFT MUD AND/OR GRAVEL, AND/O ER 1928, BUCHANAN 1980, GOODRICH AND VAN DER S		T / SOMC ER IN SMALL ST	G3 / S2S3 REAMS WHERE THE	0	0	1	0	0

Data Current as of February 2006

County Report of Endangered, Threatened, and Special Concern Plants, Animals, and Natural Communities of Kentucky Kentucky State Nature Preserves Commission

County	<b>Taxonomic Group</b>	Scientific name	Common name	Statuses	Ranks	# of Occurrences				
	Habitat					Е	Н	F	X	U
Bullitt	Freshwater Mussels INHABITS SMALL TO MEDIUM-S	Villosa lienosa SIZED RIVERS, USUALLY IN SHALLOW WATER ON A SA	Little Spectaclecase AND/MUD/DETRITUS BOTTOM (PARMALEE 1	S / 967, GORDON A	G5 / S3S4 .ND LAYZER 1989).	1	0	0	0	0
Bullitt	Crustaceans FLAT COBBLE AND BOULDER S	Orconectes jeffersoni STREWN STREAMS.	Louisville Crayfish	E/SOMC	G1 / S1	1	0	0	0	0
		Lota lota RALLY COME FROM MEDIUM TO LARGE-SIZE RIVERS. SCOTT AND CROSSMAN 1973, SMITH 1979, TRAUTMA	· · · · · · · · · · · · · · · · · · ·	S / GE, AND DEEP R	G5 / SU IVERS AND LAKES (	1	0	0	0	0
	Fishes LARGE STREAMS AND RIVERS WARREN 1986, ETNIER AND ST	Noturus stigmosus IN MODERATE TO SWIFT CURRENT OVER GRAVEL AI ARNES 1993).	Northern Madtom ND SAND, AND SOMETIMES DEBRIS OR POI	S / SOMC NDWEED FOR C	G3 / S2S3 OVER (BURR AND	2	1	0	0	0
Bullitt	Breeding Birds Open situations with scattered bus	Chondestes grammacus shes and trees, prairie, forest edge, cultivated areas, orchai	Lark Sparrow rds, fields with bushy borders, and savanna (B8	T / 3COM01NA).	G5 / S2S3B	0	1	0	0	0
Bullitt	Breeding Birds MARSHES, SWAMPS, LAKES, LA	<i>Nyctanassa violacea</i> AGOONS, AND MANGROVES.	Yellow-crowned Night-heron	Τ/	G5 / S2B	1	0	0	0	0
Bullitt	Mammals Indiana bats use primarily caves fo	Myotis sodalis or hibernacula, although they are occasionally found in old	Indiana Bat mine portals.	E/LE	G2 / S1S2	2	1	0	1	0
Bullitt	Mammals THE EVENING BAT IS A COLON	Nycticeius humeralis IAL SPECIES THAT ROOSTS IN TREES AND HOUSES.	Evening Bat IT APPARENTLY MIGRATES SOUTHWARD IN	S / N WINTER.	G5 / S3	3	0	0	0	0
Bullitt	Communities	Dolomite glade		1	GNR / S2	2	0	0	1	0
Bullitt	Communities	Limestone barrens		1	GNR / S2	1	0	0	0	0

Data Current as of February 2006 Page 5 of 5